

Anisotropic Evolution Driven by Kinetic Term

Masato Ito *

Department of Physics, Aichi University of Education, Kariya, 448-8542, JAPAN

Abstract

We present a simple model where anisotropic evolution is driven by kinetic term in extra dimensions. By introducing a canonical or a ghost kinetic term, the possibility of anisotropy is studied.

*mito@aecc.aichi-edu.ac.jp

If starting from theories such as string/M-theory or braneworld model, it is believed that extra dimensions exist at early universe [1, 2, 3, 4]. Since we live in 4-dimensional spacetime, the extra dimensions are invisible. In the string/M-theory, it is expected that extra dimensions are compactified by moduli fields and some dynamics. According to the picture of braneworld, we live in a 3-brane embedded in higher dimensional world [5]. Moreover, the Randall-Sundrum model with warped metric have shown that the fifth dimension is noncompact [6]. However, whether the extra dimensions exist isn't confirmed by experiments till now.

Recent precision of astronomical observations has illustrated whole history of time evolution of our universe [7, 8]. Accelerated expansion of universe occurs at early time and late time, the former is inflation and the latter is present cosmic acceleration due to unknown dark energy. One of the most significant problems of particle physics and cosmology is to explain the whole history of our universe. It is important to propose cosmological models to be consistent with observations. However it isn't easy to construct the consistent cosmological models from the effective theory of string/M-theory.

Assuming that the birth of our universe is in higher dimensional spacetime, we suggest that the cosmological model has anisotropic evolution of scale factors at early universe. Accordingly 3-dimensional space is expanding while extra dimensional space is contracting. In the framework of 4-dimensional spacetime, an example of vacuum cosmological model with anisotropy had been proposed by Kasner [9, 10]. The scale factor of each direction has anisotropic power-law solution t^{p_i} ($i = 1, 2, 3$), and the exponents are subject to constraints: $\sum_{i=1}^3 p_i = \sum_{i=1}^3 p_i^2 = 1$. The straightforward extension to higher dimensional spacetime of the vacuum Kasner model have been performed [11, 12, 13]. Several anisotropic models have been proposed [14, 15, 16, 17, 18, 20, 19]. We would like to consider the anisotropic evolution in extra dimension, far from the string/M-theory cosmologies. In particular, we focus on the role of kinetic term. For example it have been proposed that the kinetic term plays an important role in cosmological models such as k-inflation [21] or k-essence [22, 23, 24, 25].

In this paper we present a simple model realizing anisotropic evolution driven by kinetic term without potential. We indicate that a canonical or a ghost kinetic term leads to anisotropy. The line element with homogeneous, anisotropic and spatially flat is

$$ds^2 = -dt^2 + a^2(t) \sum_{i=1}^3 dx_i^2 + b^2(t) \sum_{M=1}^n dy_M^2, \quad (1)$$

where the scale factor of 3-dimensional space is $a(t)$ and the scale factor of extra n -

dimensional space is $b(t)$. We consider a $(4+n)$ -dimensional model with the kinetic term of a homogeneous real scalar field $\phi(t)$ without potential. The action is given by

$$S = \int d^{4+n}x \sqrt{-g} \left(\frac{1}{2} \mathcal{R} - \epsilon \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi \right), \quad (2)$$

where the higher dimensional gravitational constant is set to 1. By choosing the sign $\epsilon = \pm 1$ of kinetic term, we want to seek the possibility of anisotropic evolution. The case of $\epsilon = +1$ is canonical scalar field and the case of $\epsilon = -1$ is ghost field. Likewise, anisotropic model with ghost kinetic term without potential have been already proposed [26]. The model is constructed under the assumptions that first time derivative for Hubble parameter of each direction is zero and $\ddot{\phi} = 0$ for ghost field. In the model proposed here, the assumptions are relaxed. The present model corresponds to the combination of both [12] and [26].

In the case of choice $\epsilon = -1$, the existence of ghost field leads to the instability of physical system due to unbounded kinetic energy. Assuming that the birth of our universe comes about by instability of physical system, the ghost field can be temporarily allowed. As argued in [26], it is possible that the ghost field generates from string theory, supergravity and negative tension brane [27, 28, 29, 30].

From the setup of (1) and (2), three components of the gravitational equation and equation of motion for ϕ are given by

$$3 \left(\frac{\dot{a}}{a} \right)^2 + \frac{1}{2} n(n-1) \left(\frac{\dot{b}}{b} \right)^2 + 3n \frac{\dot{a}\dot{b}}{ab} = \epsilon \frac{1}{2} \dot{\phi}^2, \quad (3)$$

$$2 \frac{\ddot{a}}{a} + \left(\frac{\dot{a}}{a} \right)^2 + n \frac{\ddot{b}}{b} + \frac{1}{2} n(n-1) \left(\frac{\dot{b}}{b} \right)^2 + 2n \frac{\dot{a}\dot{b}}{ab} = -\epsilon \frac{1}{2} \dot{\phi}^2, \quad (4)$$

$$3 \frac{\ddot{a}}{a} + 3 \left(\frac{\dot{a}}{a} \right)^2 + (n-1) \frac{\ddot{b}}{b} + \frac{1}{2} (n-1)(n-2) \left(\frac{\dot{b}}{b} \right)^2 + 3(n-1) \frac{\dot{a}\dot{b}}{ab} = -\epsilon \frac{1}{2} \dot{\phi}^2, \quad (5)$$

$$\epsilon \left(\ddot{\phi} + \left(3 \frac{\dot{a}}{a} + n \frac{\dot{b}}{b} \right) \dot{\phi} \right) = 0, \quad (6)$$

where the dot denotes the derivative with respect to t . Here ϕ is properly normalized to be dimensionless.

We can seek the solutions to be consistent with Eqs.(3)-(6). Consequently, two scale factors $a(t)$ and $b(t)$ becomes power-law solutions or exponential-law solutions via the time-dependence of $\phi(t)$. Below we show two type of solutions separately.

Solution 1: The power-law solutions of scale factors are given by [†]

$$a(t) = a_0 t^p, \quad b(t) = b_0 t^q, \quad \phi(t) = \phi_0 \log t + \text{const}, \quad (7)$$

where a_0, b_0, ϕ_0 are real constants. Moreover two exponents p and q are subject to the following constraints

$$3p + nq = 1, \quad 3p^2 + nq^2 = 1 - \epsilon \phi_0^2. \quad (8)$$

When $\phi_0 = 0$, the model corresponds to vacuum Kasner model [11, 12, 13]. Eq.(8) yields

$$p = \frac{3 \pm \sqrt{3n(n+2 - \epsilon(n+3)\phi_0^2)}}{3(n+3)}, \quad (9)$$

$$q = \frac{n \mp \sqrt{3n(n+2 - \epsilon(n+3)\phi_0^2)}}{n(n+3)}. \quad (10)$$

Each exponent depends on n, ϕ_0 and ϵ . We investigate the anisotropy for $\epsilon = \pm 1$.

In the case of canonical scalar ($\epsilon = +1$), the reality of exponents leads to

$$0 < \phi_0^2 \leq \frac{n+2}{n+3}. \quad (11)$$

In particular, since square root is zero when $\phi_0^2 = (n+2)/(n+3)$, we obtain the isotropic evolution corresponding to $p = q = 1/(n+3)$. Thus the anisotropy depends on the value of $0 < \phi_0^2 < (n+2)/(n+3)$. Selecting the upper sign in Eqs.(9) and (10), 3-dimensional space is decelerated expansion while extra dimensional space is contracting. The value of ϕ_0 determines whether evolution is anisotropy or isotropy.

In the case of ghost ($\epsilon = -1$), the reality of exponent is automatically satisfied. For arbitrary ϕ_0 , the model has the phase of anisotropic evolution ($p \neq q$). We consider that 3-dimensional space is accelerated expansion. Since the condition is $p > 1$ (power-law inflation) for upper sign, we have

$$\phi_0^2 > 2 \frac{n+2}{n}. \quad (12)$$

On the other hand, extra dimensional space becomes accelerated contraction.

Thus, for a canonical scalar (depending on ϕ_0) or a ghost, logarithmic dependence of $\phi(t)$ leads to the anisotropic power-law solutions of scale factors.

[†]see Appendix

Solution 2: The exponential-law solutions of scale factors are given by[‡]

$$a(t) = a_0 e^{\alpha t}, \quad b(t) = b_0 e^{-3\alpha t/n}, \quad \phi(t) = \phi_0 t + \text{const}, \quad (13)$$

where ϕ_0, α are real constants and are subject to the following equation

$$3(n+3)\alpha^2 = -\epsilon n \phi_0^2. \quad (14)$$

From Eq.(14), the case of canonical scalar ($\epsilon = +1$) is excluded. In the case of ghost ($\epsilon = -1$), we have

$$\alpha = \phi_0 \sqrt{\frac{n}{3(n+3)}}, \quad (15)$$

where sign $\pm$ due to square root is absorbed in ϕ_0 .

According to Eqs.(13) and (15), positive ϕ_0 gives rise to accelerated expansion of our universe. On the other hand, extra dimensional space is accelerated contraction. Actually the solutions are described in [26]. Thus, for a ghost, time dependence of $\ddot{\phi}(t) = 0$ leads to the anisotropic exponential-law solutions of scale factors.

In conclusion we have presented a simple model with anisotropy driven by kinetic term in extra dimensions. We suggest the possibility that the kinetic term plays an important role at birth of our universe. As shown in **solution 1**, the canonical or ghost kinetic term leads to the power-law anisotropy. In the case of canonical kinetic term, 3-dimensional space is decelerated expansion and extra dimensions is accelerated contraction. When the equality holds in Eq.(11), the evolution becomes isotropy. Thus the value of ϕ_0 determines whether evolution is anisotropy or isotropy. In the case of ghost kinetic term, when inequality of Eq.(12) holds, 3-dimensional space is accelerated expansion (power-law inflation) and extra dimensions is accelerated contraction. The power-law anisotropy isn't discussed in [26]. As shown in **solution 2**, ghost kinetic term leads to the exponential-law anisotropy. The case had been described in [26].

In the case of allowing the temporarily ghost kinetic term, it is assumed that the instability of dynamics triggers the birth of our universe. We need some mechanisms in order that ghost kinetic energy is bounded. In this paper we don't discuss it.

The model we considered here may be strange scenario. The familiar early universe scenario is from inflaton-driven inflation phase due to a nearly constant potential to reheating phase at minimum of potential. We suggest that the present model is applied

[‡]see Appendix

to the birth of universe based on anisotropy of higher dimension. The mechanism of transition from anisotropic evolution to the familiar inflation is needed. Although the mechanism is unknown, we emphasised an important role of kinetic term at early universe in the paper.

Appendix: Solution 1 and 2 can be shown as follows. Integrating (6), we obtain

$$\dot{\phi} \propto a^{-3} b^{-n}. \quad (16)$$

The some combinations of Eqs.(3)-(5) lead to the following equations.

$$(n+1) \times (3) - 3 \times (4) - n \times (5) \rightarrow 3 \frac{\ddot{a}}{a} + n \frac{\ddot{b}}{b} = -\epsilon \dot{\phi}^2, \quad (17)$$

$$(3) - (n-1) \times (4) + n \times (5) \rightarrow \frac{\ddot{a}}{a} + 2 \left(\frac{\dot{a}}{a} \right)^2 + n \frac{\dot{a} \dot{b}}{a b} = 0, \quad (18)$$

$$(3) + 3 \times (4) - 2 \times (5) \rightarrow \frac{\ddot{b}}{b} + (n-1) \left(\frac{\dot{b}}{b} \right)^2 + 3 \frac{\dot{a} \dot{b}}{a b} = 0. \quad (19)$$

Using Eqs.(16)-(19), it is easy to obtain the solutions. In the case of $\dot{\phi} \propto t^{-1}$ (**solution 1**), scale factors are power-law solution. In the case of $\dot{\phi} \propto 1$ (**solution 2**), scale factors are exponential-law solution.

Acknowledgements: M.I is financially supported by Aichi University of Education.

References

- [1] M. Gasperini and G. Veneziano, *Astropart. Phys.* **1**, 317 (1993) [arXiv:hep-th/9211021].
- [2] F. Quevedo, *Class. Quant. Grav.* **19**, 5721 (2002) [arXiv:hep-th/0210292].
- [3] M. Gasperini and G. Veneziano, “String Theory and Pre-big bang Cosmology,” arXiv:hep-th/0703055.
- [4] L. McAllister and E. Silverstein, “String Cosmology: A Review,” *Gen. Rel. Grav.* **40**, 565 (2008) [arXiv:0710.2951 [hep-th]].
- [5] L. Randall and R. Sundrum, “A large mass hierarchy from a small extra dimension,” *Phys. Rev. Lett.* **83**, 3370 (1999) [arXiv:hep-ph/9905221].

- [6] L. Randall and R. Sundrum, “An alternative to compactification,” *Phys. Rev. Lett.* **83**, 4690 (1999) [arXiv:hep-th/9906064].
- [7] M. R.olta et al. [WMAP Collaboration], “Five-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Angular Power Spectra,” *Astrophys. J. Suppl.* **180**, 296 (2009) [arXiv:0803.0593 [astro-ph]].
- [8] G. Hinshaw et al. [WMAP Collaboration], “Five-Year Wilkinson Microwave Anisotropy Probe (WMAP 1) Observations: Data Processing, Sky Maps, & Basic Results,” *Astrophys. J. Suppl.* **180**, 225 (2009) [arXiv:0803.0732 [astro-ph]].
- [9] E. Kasner, *Am. J. Math* **43**, 217 (1921)
- [10] C.W. Misner, K.S. Thorne and J.A. Wheeler, *Gravitation*, (W.H. Freeman 1973)
- [11] A. Chodos and S. Detweiler, “Where Has The Fifth-Dimension Gone?,” *Phys. Rev. D* **21**, 2167 (1980).
- [12] J. J. Levin, “Inflation from extra dimensions,” *Phys. Lett. B* **343**, 69 (1995) [arXiv:gr-qc/9411041].
- [13] N. Arkani-Hamed, S. Dimopoulos, N. Kaloper and J. March-Russell, “Rapid asymmetric inflation and early cosmology in theories with sub-millimeter dimensions,” *Nucl. Phys. B* **567**, 189 (2000) [arXiv:hep-ph/9903224].
- [14] J. M. Cline, “Inflation from extra dimensions,” *Phys. Rev. D* **61**, 023513 (2000) [arXiv:hep-ph/9904495].
- [15] A. Mazumdar, “Extra dimensions and inflation,” *Phys. Lett. B* **469**, 55 (1999) [arXiv:hep-ph/9902381].
- [16] A. Feinstein and S. Jhingan, “Ghosts in a mirror,” *Mod. Phys. Lett. A* **19**, 457 (2004) [arXiv:hep-th/0304069].
- [17] S. Chatterjee, A. Banerjee and Y. Z. Zhang, “Accelerating universe from extra spatial dimension,” *Int. J. Mod. Phys. A* **21**, 4035 (2006) [arXiv:gr-qc/0509112].
- [18] B. R. Greene and J. Levin, “Dark Energy and Stabilization of Extra Dimensions,” *JHEP* **0711**, 096 (2007) [arXiv:0707.1062 [hep-th]].

- [19] T. Koivisto and D. F. Mota, “Accelerating Cosmologies with an Anisotropic Equation of State,” *Astrophys. J.* **679**, 1 (2008) [arXiv:0707.0279 [astro-ph]].
- [20] T. S. Koivisto and D. F. Mota, “Vector Field Models of Inflation and Dark Energy,” *JCAP* **0808**, 021 (2008) [arXiv:0805.4229 [astro-ph]].
- [21] C. Armendariz-Picon, T. Damour and V. F. Mukhanov, “k-Inflation,” *Phys. Lett. B* **458**, 209 (1999) [arXiv:hep-th/9904075].
- [22] T. Chiba, T. Okabe and M. Yamaguchi, “Kinetically driven quintessence,” *Phys. Rev. D* **62**, 023511 (2000) [arXiv:astro-ph/9912463].
- [23] C. Armendariz-Picon, V. F. Mukhanov and P. J. Steinhardt, “Essentials of k-essence,” *Phys. Rev. D* **63**, 103510 (2001) [arXiv:astro-ph/0006373].
- [24] T. Chiba, “Tracking k-essence,” *Phys. Rev. D* **66**, 063514 (2002) [arXiv:astro-ph/0206298].
- [25] L. P. Chimento and A. Feinstein, “Power-law expansion in k-essence cosmology,” *Mod. Phys. Lett. A* **19**, 761 (2004) [arXiv:astro-ph/0305007].
- [26] S. P. Patil, “Exponential inflation with $\rho = +p$,” arXiv:hep-th/0509088.
- [27] I. I. Kogan, S. Mouslopoulos, A. Papazoglou and L. Pilo, “Radion in Multibrane World,” *Nucl. Phys. B* **625**, 179 (2002) [arXiv:hep-th/0105255].
- [28] P. L. McFadden and N. Turok, “Conformal symmetry of brane world effective actions,” *Phys. Rev. D* **71**, 021901 (2005) [arXiv:hep-th/0409122].
- [29] I. Y. Aref’eva, A. S. Koshelev and S. Y. Vernov, “Exactly Solvable SFT Inspired Phantom Model,” *Theor. Math. Phys.* **148**, 895 (2006) [*Teor. Mat. Fiz.* **148**, 23 (2006)] [arXiv:astro-ph/0412619].
- [30] I. Y. Aref’eva, “Nonlocal string tachyon as a model for cosmological dark energy,” *AIP Conf. Proc.* **826**, 301 (2006) [arXiv:astro-ph/0410443].